D-192 袖珍椰子. Parlor palm. chamaedorea elegans Mart.

O_3 0.508 ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後2天.

O_3 0.508 ppm, fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃ (day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 2 days after fumigation.

D-193 袖珍椰子. Parlor palm.

再觀察照片 D-192. (攝影：處理後 4天). (徵狀：如照片所示).

Again a view of plate D-192. (photo: 4 days after fumigation). (symptom:as shown in the plate).

D-194 袖珍椰子. Parlor palm.

再觀察照片 D-192. (攝影：處理後 6天). (徵狀：如照片所示).

Again a view of plate D-192. (photo: 6 days after fumigation). (symptom:as shown in the plate).

D-195 相思樹. Acacia. Acacia sp.

O_3 0.82ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80%，光度7000lux，徵狀：如照片所示，攝影：處理後1天，使用品種：台灣本地種.

O_3 0.82ppm, fumigation period: 24 hrs. (day/night), fumigation condition: temp. 25／20℃ (day/night), humidity 80%, light 7000 lux, symptom: as shown in the plate, photo: 1 day after fumigation, cultivar used is written in the Chinese section.

D-196 相思樹. Acacia.

照片 D-195 的近照. (徵狀：如照片所示).

A close view of plate D-195. (symptom:as shown in the plate).



D-197 相思樹. Acacia.

再觀察照片 D-195. (攝影：處理後 7天). (徵狀：如照片所示).

Again a view of plate D-195. (photo: 7 days after fumigation). (symptom:as shown in the plate).

D-198 相思樹. Acacia.

照片 D-197 的近照. (徵狀：如照片所示).

A close view of plate D-197. (symptom:as shown in the plate).

D-199 腎蕨.

Erect sword fern. Nephrolepis cordifolia K. B. PRESL.

O_3 0.6 ppm，處理時間：1小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後1天.

O_3 0.6 ppm, fumigation period: 1 hr. (day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 1 day after fumigation.

E. 混合氣體污染物. Air Pollutant Combinations.

(二氧化硫＋二氧化氮) (Sulfur Dioxide ＋ Nitrogen Dioxide)

E-1 蕃石榴. Guava. Psidium guajava L.

SO_2 0.48 ppm + NO_2 3.1 ppm，處理時間：5天(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：葉尖乾枯及葉緣乾枯，攝影：處理後7天，使用品種：梨拔.

SO_2 0.48 ppm + NO_2 3.1 ppm, fumigation period: 5 days.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: tip necrosis & marginal necrosis, photo: 7 days after fumigation, cultivar used is written in the Chinese section.

E-2 蕃石榴. Guava.

照片E-1的近照.(葉尖乾枯及葉緣乾枯).

A close view of plate E-1. (tip necrosis & marginal necrosis).

E-3 蕃石榴. Guava.

再觀察照片 E-1. (攝影：處理後17天). (葉尖乾枯及葉緣乾枯).

Again a view of plate E-1. (photo: 17 days after fumigation). (tip necrosis & marginal necrosis).

E-4 蕃石榴. Guava.

照片 E-3 的近照. (葉尖乾枯及葉緣乾枯).

A close view of plate E-3. (tip necrosis & marginal necrosis).

E-5 香蕉. Banana. Musa spp.

SO_2 0.44 ppm + NO_2 2.9 ppm，處理時間：4天(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80%，光度7000lux，徵狀：葉間水漬狀，攝影：處理後立即拍照，使用品種：北蕉.

SO_2 0.44 ppm + NO_2 2.9 ppm，fumigation period: 4 days.(day/night), fumigation condition: temp. 25／20℃ (day/night), humidity 80%, light 7000 lux, symptom: watersoaked appearance, photo: immediately after fumigation, cultivar used is written in the Chinese section.

E-6 香蕉. Banana.

再觀察照片 E-5. (攝影：處理後 8天). (葉緣乾枯).

Again a view of plate E-5. (photo: 8 days after fumigation). (marginal necrosis).

E-7 香蕉. Banana.

SO_2 0.48ppm + NO_2 3.1ppm，處理時間：5天(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：葉緣乾枯，攝影：處理後17天，使用品種：北蕉.

SO_2 0.48ppm + NO_2 3.1ppm, fumigation period: 5 days.(day/night), fumigation condition: temp. 25 ／ 20 ℃ (day/night), humidity 80％, light 7000 lux, symptom: marginal necrosis, photo: 17 days after fumigation, cultivar used is written in the Chinese section.

E-8 相思樹. Acacia. Acacia sp.

SO_2 0.29ppm + NO_2 1.9ppm，處理時間：4天(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示(右邊葉)，攝影：處理後9天，使用品種：台灣本地種.

SO_2 0.29ppm + NO_2 1.9ppm，fumigation period: 4 days.(day/night), fumigation ondition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate (right hand leaf), photo: 9 days after fumigation, cultivar used is written in the Chinese section.

(二氧化硫＋氟化物) (Sulfur Dioxide ＋ Fluorides)

E-9 蕃石榴. Guava. Psidium guajava L.

SO_2 2.8 ppm + HF 0.08ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000 lux，徵狀：葉間水漬狀，攝影：處理後立即拍照，使用品種：梨拔.

SO_2 2.8 ppm + HF 0.08ppm, fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: watersoaked appearance, photo: immediately after fumigation, cultivar used is written in the Chinese section.

E-10 蕃石榴. Guava.

再觀察照片 E-9. (攝影：處理後2天). (如照片所示的葉間乾枯).

Again a view of plate E-9.
(photo: 2 days after fumigation).
(necrosis as shown in the plate).

E-11 蕃石榴. Guava.

再觀察照片 E-9. (攝影：處理後2天). (如照片所示的葉間乾枯).

Again a view of plate E-9.
(photo: 2 days after fumigation).
(necrosis as shown in the plate).

E-12 橄欖. Chinese olive. Canarium album Raeusch.

SO_2 2.8 ppm + HF 0.08ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後9天，使用品種：台灣本地種.

SO_2 2.8 ppm + HF 0.08 ppm, fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃ (day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 9 days after fumigation, cultivar used is written in the Chinese section.

E-13 橄欖. Chinese olive.

照片 E-12 的另一角度攝影.(徵狀：如照片所示).

Another view of plate E-12.
(symptom:as shown in the plate).

E-14 唐菖蒲. Gladioli. Gladiolus hybridus Hort.



SO_2 2.8 ppm + HF 0.08ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後立即拍照，使用品種：新粉.

SO_2 2.8 ppm + HF 0.08ppm, fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／ 20℃(day/night), humidity 80%, light 7000 lux, symptom: as shown in the plate, photo: immediately after fumigation, cultivar used is written in the Chinese section.



E-15 唐菖蒲. Gladioli.

再觀察照片 E-14.(攝影：處理後2天).(徵狀：如照片所示).

Again a view of plate E-14.
(photo: 2 days after fumigation). (symptom: as shown in the plate).

E-16 唐菖蒲. Gladioli.

再觀察照片 E-14. (攝影：處理後 9 天). (徵狀：如照片所示).

Again a view of plate E-14. (photo: 9 days after fumigation). (symptom:as shown in the plate).

E-17 相思樹. Acacia. Acacia sp.

SO_2 2.8 ppm + HF 0.08ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後立即拍照，使用品種：台灣本地種.

SO_2 2.8 ppm + HF 0.08 ppm, fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃ (day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: immediately after fumigation, cultivar used is written in the Chinese section.

E-18 相思樹. Acacia.

再觀察照片 E-17. (攝影：處理後 2天). (徵狀：如照片所示).
Again a view of plate E-17. (photo: 2 days after fumigation). (symptom:as shown in the plate).

E-19 相思樹. Acacia.

再觀察照片 E-17.(攝影：處理後9天).(徵狀：如照片所示).

Again a view of plate E-17. (photo: 9 days after fumigation). (symptom:as shown in the plate).

(臭氧＋氟化物) (Ozone + Fluorides)

E-20 木薯. Cassava. Manihot esculenta Crantz.

O_3 0.33ppm + HF 0.32ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000 lux，徵狀：如照片所示，攝影：處理後2天，使用品種：甜味種.

O_3 0.33ppm + HF 0.32ppm, fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 2 days after fumigation, cultivar used is written in the Chinese section.

E-21 菸草. Tobacco. Nicotiana tabacum L.

O_3 0.2 ppm + HF 0.24ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後3天，使用品種：台菸6號.

O_3 0.2 ppm + HF 0.24ppm, fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20 ℃ (day/night), humidity 80 %, light 7000 lux, symptom: as shown in the plate, photo: 3 days after fumigation, cultivar used is written in the Chinese section.

E-22 菸草. Tobacco.

照片 E-21 的近照. (徵狀：如照片所示).

A close view of plate E-21.
(symptom: as shown in the plate).



E-23 菸草. Tobacco.

照片 E-21 的近照. (徵狀：如照片所示).

A close view of plate E-21.
(symptom: as shown in the plate).

E-24 香蕉. Banana. Musa spp.

O_3 0.3 ppm + HF 0.32ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後8天，使用品種：Willian Hybrid.

O_3 0.3 ppm + HF 0.32 ppm，fumigation period: 24 hrs. (day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 8 days after fumigation, cultivar used is written in the Chinese section.

E-25 香蕉. Banana.

O_3 0.3 ppm + HF 0.32ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000 lux，徵狀：如照片所示，攝影：處理後8天，使用品種：215.

O_3 0.3 ppm + HF 0.32ppm，fumigation period: 24 hrs. (day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 8 days after fumigation, cultivar used is written in the Chinese section.

E-26 香蕉. Banana.

O_3 0.3 ppm + HF 0.32ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後8天，使用品種：Valery.

O_3 0.3 ppm + HF 0.32ppm，fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 8 days after fumigation, cultivar used is written in the Chinese section.

E-27 咖啡. Coffee. Coffea spp.

O_3 0.33ppm + HF 0.32ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後立即拍照，使用品種：適合台灣栽培品系.

O_3 0.33ppm + HF 0.32ppm，fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: immediately after fumigation, cultivar used is written in the Chinese section.

E-28 蓮霧. Wax apple. Syzygium samarangense Merr. et perry.

O_3 0.33ppm + HF 0.32ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後立即拍照，使用品種：黑珍珠.

O_3 0.33ppm + HF 0.32ppm，fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: immediately after fumigation, cultivar used is written in the Chinese section.

E-29 木瓜. Papaya. Carica papaya Linn.

O_3 0.3 ppm + HF 0.32ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度 80％，光度 7000 lux，徵狀：葉間水漬狀，攝影：處理後 1天，使用品種：日陞.

O_3 0.3 ppm + HF 0.32ppm，fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: watersoaked appearance, photo: 1 day after fumigation, cultivar used is written in the Chinese section.

E-30 木瓜. Papaya.



再觀察照片 E-29. (攝影：處理後8天). (葉面乾枯及落葉).

Again a view of plate E-29. (photo: 8 days after fumigation). (necrosis & defoliation).

E-31 菜. Water spinach. Ipomoea aquatica Forsk.

O_3 0.2 ppm + HF 0.24ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後1天，使用品種：大葉種.

O_3 0.2 ppm + HF 0.24ppm，fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 1 day after fumigation, cultivar used is written in the Chinese section.

E-32 菜. Water spinach.

照片 E-31 的近照. (徵狀：如照片所示).

A close view of plate E-31.
(symptom: as shown in the plate).

E-33 茄子. Egg-plant. Solanum melongena L.



O_3 0.2 ppm + HF 0.24ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後1天，使用品種：豐原 薯茄.

O_3 0.2 ppm + HF 0.24ppm，fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 1 day after fumigation, cultivar used is written in the Chinese section.

E-34 茄子. Egg-plant.



照片 E-33 的近照.(徵狀：如照片所示).

A close view of plate E-33. (symptom: as shown in the plate).

E-35 茄子. Egg-plant.



再觀察照片 E-34.(攝影：處理後3天).(徵狀：如照片所示).

Again a view of plate E-34. (photo: 3 days after fumigation). (symptom:as shown in the plate).

E-36 黃秋葵. Okra. Hibiscus esculentus L.



O_3 0.2 ppm + HF 0.24ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後1天，使用品種：農友五福.

O_3 0.2 ppm + HF 0.24ppm，fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃(day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 1 day after fumigation, cultivar used is written in the Chinese section.

E-37 黃秋葵. Okra.



照片E-36的近照. (受害不嚴重之葉). (徵狀：如照片所示).

A close view of plate E-36. (not severely injured leaf). (symptom: as shown in the plate).

E-38 黃秋葵. Okra.



再觀察照片E-37. (攝影：處理後3天). (徵狀：如照片所示).

Again a view of plate E-37. (photo: 3 days after fumigation). (symptom:as shown in the plate).

E-39 毛豆. Vegetable soybean. Glycine Max Merr.



O_3 0.33ppm + HF 0.32ppm，處理時間：6小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80%，光度7000lux，徵狀：如照片所示，攝影：處理後2天，使用品種：台農品系.

O_3 0.33ppm + HF 0.32ppm, fumigation period: 6 hrs.(day/night), fumigation condition: temp. 25／20℃ (day/night), humidity 80%, light 7000 lux, symptom: as shown in the plate, photo: 2 days after fumigation, cultivar used is written in the Chinese section.

E-40 毛豆. Vegetable soybean.

照片 E-39 的近照. (受害嚴重之葉). (如照片所示的葉間乾枯).

A close view of plate E-39.
(severely injured leaves).
(necrosis as shown in the plate).

E-41 毛豆. Vegetable soybean.

照片 E-39 的近照. (受害不嚴重之葉). (徵狀：如照片所示).

A close view of plate E-39.
(not severely injured leaf).
(symptom: as shown in the plate).

E-42 金露花. Duranta repens.

O_3 0.2 ppm + HF 0.24ppm，處理時間：24小時(日／夜)，處理時條件：溫度25／20℃(日／夜)，溼度80％，光度7000lux，徵狀：如照片所示，攝影：處理後3天.

O_3 0.2 ppm + HF 0.24ppm，fumigation period: 24 hrs.(day/night), fumigation condition: temp. 25／20℃ (day/night), humidity 80％, light 7000 lux, symptom: as shown in the plate, photo: 3 days after fumigation.

E-43 金露花. Duranta repens.



照片 E-42 的近照.(徵狀：如照片所示).

A close view of plate E-42. (symptom: as shown in the plate).

E-44 金露花. Duranta repens.

照片 E-42 的近照.(徵狀：如照片所示).

A close view of plate E-42. (symptom: as shown in the plate).

索 引 (Index)

Page.

A.二氧化硫・Sulfur Dioxide.

1	A-1	水稻・Rice. Oryza sativa L.
1	A-2	水稻・Rice.
2	A-3	水稻・Rice.
2	A-4	水稻・Rice.
3	A-5	水稻・Rice.
3	A-6	甘藷・Sweet Potato. Ipomoea batatas (L.).
4	A-7	甘藷・Sweet Potato.
4	A-8	花生・Peanut. Arachis hypogaea L.
5	A-9	花生・Peanut.
5	A-10	大豆・Soybean. Glycine max(L.) Merrill.
6	A-11	大豆・Soybean.
6	A-12	玉米・Corn. Zea mays Linn.
7	A-13	玉米・Corn.
7	A-14	綠豆・Mungbean. Vigna radiata (L) Wilczek.
8	A-15	綠豆・Mungbean.
8	A-16	綠豆・Mungbean.
9	A-17	甘蔗・Sugarcane. Saccharum officinarum L.
9	A-18	甘蔗・Sugarcane.
10	A-19	甘蔗・Sugarcane.
10	A-20	香蕉・Banana. Musa spp.
11	A-21	香蕉・Banana.
11	A-22	香蕉・Banana.
12	A-23	鳳梨・Pineapple. Ananas comosus Merr.
12	A-24	龍眼・Longan. Euphoria longana Lam.
13	A-25	蕃石榴・Guava. Psidium guajava L.
13	A-26	蕃石榴・Guava.
14	A-27	橄欖・Kanran (Chinese olive). Canarium album Raeusch.
14	A-28	橄欖・Kanran (Chinese olive).
15	A-29	橄欖・Kanran (Chinese olive).
15	A-30	橄欖・Kanran (Chinese olive).
16	A-31	咖啡・Coffee. Coffea spp.

16 A-32 咖啡・Coffee.
17 A-33 柑桔・Citrus fruit. citrus spp.
17 A-34 蓮霧・Wax apple. Syzygium samarangense Merr.et Perry.
18 A-35 芒果・Mango. Mangifera indica L.
18 A-36 芒果・Mango.
19 A-37 芒果・Mango.
19 A-38 木瓜・Papaya. Carica papaya Linn.
20 A-39 木瓜・Papaya.
20 A-40 可可椰子・Coconut. Cocos nucifera L.
21 A-41 可可椰子・Coconut.
21 A-42 檳榔・Betel palm. Areca cathecu L.
22 A-43 楊桃・Carambola. Averrhoa carambola, L.
22 A-44 楊桃・Carambola.
23 A-45 酪梨・Avocado. Persea americana Mill.
23 A-46 酪梨・Avocado.
24 A-47 酪梨・Avocado.
24 A-48 波羅蜜・Jack fruit. Artocarpus heterophyllus Lam.
25 A-49 波羅蜜・Jack fruit.
25 A-50 錫蘭橄欖・Ceylon olive. Elaeocartus Serratus L.
26 A-51 梨・Pear. Pyrus communis L.
26 A-52 梨・Pear.
27 A-53 梨・Pear.
27 A-54 梨・Pear.
28 A-55 葡萄・Grape. Vitis vinifera L.
28 A-56 葡萄・Grape.
29 A-57 葡萄・Grape.
29 A-58 蘿蔔・Radish. Raphanus sativus L.
30 A-59 蘿蔔・Radish.
30 A-60 甘藍・Cabbage. Brassica oleracea var. capitata L.
31 A-61 甘藍・Cabbage.
31 A-62 萵苣・Lettuce. Lactuca sativa L.
32 A-63 萵苣・Lettuce.
32 A-64 芥菜・Mustard. Brassica juncea Coss.
33 A-65 芥菜・Mustard.
33 A-66 芥菜・Mustard.

34 A-67 蕹菜・Water spinach. Ipomoea aquatica Forsk.
34 A-68 蕹菜・Water spinach.
35 A-69 蕹菜・Water spinach.
35 A-70 蔥 ・Spring onion. Allium fistulosum L.
36 A-71 蔥 ・Spring onion.
36 A-72 結球白菜・Chinese heading cabbage. Brassica pekinensis Rupr.
37 A-73 結球白菜・Chinese heading cabbage.
37 A-74 結球白菜・Chinese heading cabbage.
38 A-75 芥藍・Kale. Brassica oleracea L. var. acephala DC.
38 A-76 芥藍・Kale.
39 A-77 芥藍・Kale.
39 A-78 花椰菜・Cauliflower. Brassica oleracea L. var. botrytis L.
40 A-79 花椰菜・Cauliflower.
40 A-80 黃瓜・Cucumber. Cucumis sativus L.
41 A-81 黃瓜・Cucumber.
41 A-82 黃瓜・Cucumber.
41 A-83 絲瓜・Vegetable sponge. Luffa spp.
42 A-84 絲瓜・Vegetable sponge.
42 A-85 南瓜・Pumpkin. Cucurbita sp.
43 A-86 冬瓜・Wax gourd. Benincasa hispida Cogn.
43 A-87 冬瓜・Wax gourd.
44 A-88 苦瓜・Bitter gourd. Momordica charantia L.
44 A-89 苦瓜・Bitter gourd.
44 A-90 苦瓜・Bitter gourd.
45 A-91 甜椒・Sweet pepper. Capsicum annuum L.
45 A-92 甜椒・Sweet pepper.
46 A-93 豇豆・Yard long bean. Vigna sesquipedalis Fruwirth.
46 A-94 豇豆・Yard long bean.
46 A-95 豇豆・Yard long bean.
47 A-96 扁蒲・Bottle gourd. Lagenaria siceraria Standl.
47 A-97 扁蒲・Bottle gourd.
48 A-98 扁蒲・Bottle gourd.
48 A-99 菊花・Floret chrysanthemum. Chrysanthemum morifolium Ramat.
49 A-100 菊花・Floret chrysanthemum.
49 A-101 唐菖蒲・Gladioli. Gladiolus hybridus Hort.

50 A-102 唐菖蒲・Gladioli.
50 A-103 唐菖蒲・Gladioli.
51 A-104 唐菖蒲・Gladioli.
51 A-105 腎蕨・Erect sword fern. Nephrolepis cordifolia K. B. PRESL.
51 A-106 腎蕨・Erect sword fern.
52 A-107 滿江紅・Azolla.
52 A-108 滿江紅・Azolla.
53 A-109 滿江紅・Azolla.
53 A-110 苜蓿・Alfalfa. Medicago sativa L.
54 A-111 苜蓿・Alfalfa.
54 A-112 彩葉草・Painted nettle. Coleus blumei BENTH.
54 A-113 彩葉草・Painted nettle.
55 A-114 彩葉芋・Fancy-leaved caladium. Caladium X hortulanum BIRDSEY.
55 A-115 彩葉芋・Fancy-leaved caladium.
56 A-116 落地生根・Air-plant. Kalanchoe pinnata Pers.
56 A-117 落地生根・Air-plant.
57 A-118 百慕達草・Bermuda-grass. Cynodon Dactylon Pers.
57 A-119 百慕達草・Bermuda-grass.
58 A-120 相思樹・Acacia. Acacia sp.
58 A-121 相思樹・Acacia.
59 A-122 相思樹・Acacia.

B. 氟化物・Fluorides.

60 B-1 水稻・Rice. Oryza sativa L.
60 B-2 水稻・Rice.
61 B-3 水稻・Rice.
61 B-4 玉米・Corn. Zea mays Linn.
62 B-5 玉米・Corn.
62 B-6 玉米・Corn.
62 B-7 玉米・Corn.
63 B-8 香蕉・Banana. Musa spp.
63 B-9 香蕉・Banana.
64 B-10 香蕉・Banana.
64 B-11 香蕉・Banana.
64 B-12 香蕉・Banana.

65	B-13	鳳梨・Pineapple. Ananas Comosus Merr.
65	B-14	鳳梨・Pineapple.
66	B-15	橄欖・Chinese olive. Canarium album Raeusch.
66	B-16	橄欖・Chinese olive.
67	B-17	橄欖・Chinese olive.
67	B-18	咖啡・Coffee. Coffea spp.
68	B-19	咖啡・Coffee.
68	B-20	可可椰子・Coconut. Cocos nucifera L.
68	B-21	可可椰子・Coconut.
69	B-22	可可椰子・Coconut.
69	B-23	可可椰子・Coconut.
69	B-24	楊桃・Carambola. Averrhoa carambola, L.
70	B-25	錫蘭橄欖・Ceylon olive. Elaeocartus Serratus L.
70	B-26	錫蘭橄欖・Ceylon olive.
70	B-27	錫蘭橄欖・Ceylon olive.
71	B-28	錫蘭橄欖・Ceylon olive.
71	B-29	錫蘭橄欖・Ceylon olive.
71	B-30	錫蘭橄欖・Ceylon olive.
72	B-31	桃・Peach. Amygdalus persica Sieb. et Zucc.
72	B-32	桃・Peach.
72	B-33	桃・Peach.
73	B-34	葡萄・Grape. Vitis Vinifera L.
73	B-35	葡萄・Grape.
74	B-36	蕹菜・Water spinach. Ipomoea aquatica Forsk.
74	B-37	蕹菜・Water spinach.
75	B-38	蕹菜・Water spinach.
75	B-39	蕹菜・Water spinach.
75	B-40	蕹菜・Water spinach.
76	B-41	蕹菜・Water spinach.
76	B-42	冬瓜・Wax gourd. Benincasa hispida Cogn.
77	B-43	冬瓜・Wax gourd.
77	B-44	唐菖蒲・Gladioli. Gladiolus hybridus Hort.
77	B-45	唐菖蒲・Gladioli.
78	B-46	唐菖蒲・Gladioli.
78	B-47	唐菖蒲・Gladioli.

79 B-48 唐菖蒲・Gladioli.
79 B-49 腎蕨・Erect sword fern. Nephrolepis cordifolia K. B. PRESL.
80 B-50 腎蕨・Erect sword fern.
80 B-51 腎蕨・Erect sword fern.
80 B-52 腎蕨・Erect sword fern.
81 B-53 腎蕨・Erect sword fern.
81 B-54 相思樹・Acacia. Acacia sp.
82 B-55 相思樹・Acacia.
82 B-56 相思樹・Acacia.

C. 氮氧化物・Oxides of nitrogen.

83 C-1 花生・Peanut. Arachis hypogaea L.
83 C-2 花生・Peanut.
84 C-3 香蕉・Banana. Musa spp.
84 C-4 香蕉・Banana.
85 C-5 香蕉・Banana.
85 C-6 香蕉・Banana.
85 C-7 香蕉・Banana.
86 C-8 咖啡・Coffee. Coffea spp.
86 C-9 咖啡・Coffee.
86 C-10 咖啡・Coffee.
87 C-11 咖啡・Coffee.
87 C-12 咖啡・Coffee.
87 C-13 波羅蜜・Jack fruit. Artocarpus heterophyllus Lam.
88 C-14 波羅蜜・Jack fruit.
88 C-15 波羅蜜・Jack fruit.
88 C-16 波羅蜜・Jack fruit.
89 C-17 甘藍・Cabbage. Brassica oleracea var. capitata L.
89 C-18 甘藍・Cabbage.
89 C-19 甘藍・Cabbage.
90 C-20 甘藍・Cabbage.
90 C-21 甘藍・Cabbage.
90 C-22 蕹菜・Water spinach. Ipomoea aquatica Forsk.
91 C-23 結球白菜・Chinese heading cabbage. Brassica pekinensis Rupr.
91 C-24 結球白菜・Chinese heading cabbage.
92 C-25 結球白菜・Chinese heading cabbage.

92 C-26 結球白菜・Chinese heading cabbage.
92 C-27 結球白菜・Chinese heading cabbage.
93 C-28 絲瓜・Vegetable sponge. Luffa spp.
93 C-29 豇豆・Yard long bean. Vigna sesquipedalis Fruwirth.
94 C-30 豇豆・Yard long bean.
94 C-31 菜豆・String bean. phaseolus vulgaris L.
94 C-32 菜豆・String bean.
95 C-33 唐菖蒲・Gladioli. Gladiolus hybridus Hort.
95 C-34 唐菖蒲・Gladioli.
96 C-35 唐菖蒲・Gladioli.
96 C-36 相思樹・Acacia. Acacia sp.
97 C-37 相思樹・Acacia.
97 C-38 相思樹・Acacia.
97 C-39 相思樹・Acacia.
98 C-40 相思樹・Acacia.
98 C-41 腎蕨・Erect sword fern. Nephrolepis cordifolia K. B. PRESL.
99 C-42 滿江紅・Azolla.

D. 臭氧・Ozone.

100 D-1 甘藷・Sweet potato. Ipomoea batatas (L.)
100 D-2 甘藷・Sweet potato.
101 D-3 甘藷・Sweet potato.
101 D-4 大豆・Soybean. Glycine max (L.) Merrill.
102 D-5 大豆・Soybean.
102 D-6 大豆・Soybean.
102 D-7 大豆・Soybean.
103 D-8 大豆・Soybean.
103 D-9 大豆・Soybean.
104 D-10 玉米・Corn. Zea mays Linn.
104 D-11 玉米・Corn.
105 D-12 玉米・Corn.
105 D-13 小麥・Wheat. Triticum vulgare Vill.
105 D-14 小麥・Wheat.
106 D-15 小麥・Wheat.
106 D-16 綠豆・Mung bean. Vigna radiata (L) Wilczek.
106 D-17 綠豆・Mung bean.

107	D-18	綠豆・Mung bean.
107	D-19	綠豆・Mung bean.
107	D-20	綠豆・Mung bean.
108	D-21	綠豆・Mung bean.
108	D-22	綠豆・Mung bean.
109	D-23	綠豆・Mung bean.
109	D-24	木薯・Cassava. Manihot esculenta Crantz.
110	D-25	木薯・Cassava.
110	D-26	木薯・Cassava.
110	D-27	木薯・Cassava.
111	D-28	木薯・Cassava.
111	D-29	木薯・Cassava.
111	D-30	木薯・Cassava.
112	D-31	木薯・Cassava.
112	D-32	甘蔗・Sugarcane. Saccharum officinarum L.
112	D-33	甘蔗・Sugarcane.
113	D-34	甘蔗・Sugarcane.
113	D-35	甘蔗・Sugarcane.
113	D-36	茶葉・Tea. Thea sinensis (L) Sims.
114	D-37	茶葉・Tea.
114	D-38	茶葉・Tea.
114	D-39	茶葉・Tea.
115	D-40	菸草・Tobacco. Nicotiana tabacum L.
115	D-41	菸草・Tobacco.
116	D-42	菸草・Tobacco.
116	D-43	菸草・Tobacco.
117	D-44	菸草・Tobacco.
117	D-45	菸草・Tobacco.
118	D-46	苜蓿・Alfalfa. Medicago sativa L.
118	D-47	苜蓿・Alfalfa.
118	D-48	苜蓿・Alfalfa.
119	D-49	香蕉・Banana. Musa spp.
119	D-50	香蕉・Banana.
120	D-51	香蕉・Banana.
120	D-52	香蕉・Banana.

121	D-53	柑桔・Citrus fruit. Citrus spp.
121	D-54	荔枝・Litchi. Litchi chinensis Sonn.
122	D-55	龍眼・Longan. Euphoria longana Lam.
122	D-56	龍眼・Longan.
122	D-57	龍眼・Longan.
123	D-58	龍眼・Longan.
123	D-59	龍眼・Longan.
123	D-60	龍眼・Longan.
124	D-61	蕃石榴・Guava. Psidium guajava L.
124	D-62	蕃石榴・Guava.
124	D-63	蕃石榴・Guava.
125	D-64	蕃石榴・Guava.
125	D-65	蕃石榴・Guava.
125	D-66	蕃石榴・Guava.
126	D-67	橄欖・Chinese olive. Canarium album Raeusch.
126	D-68	橄欖・Chinese olive.
126	D-69	橄欖・Chinese olive.
127	D-70	蕃荔枝・Sugar apple. Anona squamosa, L.
127	D-71	蕃荔枝・Sugar apple.
127	D-72	蕃荔枝・Sugar apple.
128	D-73	蕃荔枝・Sugar apple.
128	D-74	蕃荔枝・Sugar apple.
128	D-75	蕃荔枝・Sugar apple.
129	D-76	蕃荔枝・Sugar apple.
129	D-77	馬拉巴栗・Malabar chestnut. Pachira macrocarpa Schlecht.
129	D-78	馬拉巴栗・Malabar chestnut.
130	D-79	馬拉巴栗・Malabar chestnut.
130	D-80	馬拉巴栗・Malabar chestnut.
130	D-81	馬拉巴栗・Malabar chestnut.
131	D-82	馬拉巴栗・Malabar chestnut.
131	D-83	蓮霧・Wax apple. Syzygium samarangense Merr. et Perry.
132	D-84	蓮霧・Wax apple.
132	D-85	芒果・Mango. Mangifera indica L.
132	D-86	芒果・Mango.
133	D-87	芒果・Mango.

133	D-88	芒果・Mango.
133	D-89	芒果・Mango.
134	D-90	芒果・Mango.
134	D-91	檳榔・Betel palm. Areca cathecu L.
134	D-92	檳榔・Betel palm.
135	D-93	酪梨・Avocado. Persea americana Mill..
135	D-94	酪梨・Avocado.
135	D-95	酪梨・Avocado.
136	D-96	波羅蜜・Jack fruit. Artocarpus heterophyllus Lam.
136	D-97	波羅蜜・Jack fruit.
137	D-98	波羅蜜・Jack fruit.
137	D-99	波羅蜜・Jack fruit.
137	D-100	波羅蜜・Jack fruit.
138	D-101	波羅蜜・Jack fruit.
138	D-102	波羅蜜・Jack fruit.
138	D-103	錫蘭橄欖・Ceylon olive. Elaeocartus Serratus L.
139	D-104	錫蘭橄欖・Ceylon olive.
139	D-105	錫蘭橄欖・Ceylon olive.
139	D-106	錫蘭橄欖・Ceylon olive.
140	D-107	錫蘭橄欖・Ceylon olive.
140	D-108	葡萄・Grape. Vitis vinifera L.
140	D-109	葡萄・Grape.
141	D-110	葡萄・Grape.
141	D-111	葡萄・Grape.
141	D-112	草梅・Strawberry. Fragaria ananassa Duch.
142	D-113	芋・Taro. Colocasia esculenta schott.
142	D-114	芋・Taro.
143	D-115	茭白・Water bamboo. Zizania aquatica L.
143	D-116	茭白・Water bamboo.
144	D-117	馬鈴薯・Potato. Solanum tuberosum L.
144	D-118	馬鈴薯・Potato.
145	D-119	馬鈴薯・Potato.
145	D-120	馬鈴薯・Potato.
146	D-121	馬鈴薯・Potato.
146	D-122	甘藍・Cabbage. Brassica oleracea var. capitata L.

146 D-123 甘藍・Cabbage.
147 D-124 甘藍・Cabbage.
147 D-125 不結球白菜・Non-heading Chinese cabbage. Brassica sp.
147 D-126 不結球白菜・Non-heading Chinese cabbage.
148 D-127 不結球白菜・Non-heading Chinese cabbage.
148 D-128 不結球白菜・Non-heading Chinese cabbage.
148 D-129 不結球白菜・Non-heading Chinese cabbage.
149 D-130 不結球白菜・Non-heading Chinese cabbage.
149 D-131 不結球白菜・Non-heading Chinese cabbage.
149 D-132 不結球白菜・Non-heading Chinese cabbage.
150 D-133 不結球白菜・Non-heading Chinese cabbage.
150 D-134 不結球白菜・Non-heading Chinese cabbage.
150 D-135 不結球白菜・Non-heading Chinese cabbage.
151 D-136 不結球白菜・Non-heading Chinese cabbage.
151 D-137 不結球白菜・Non-heading Chinese cabbage.
151 D-138 不結球白菜・Non-heading Chinese cabbage.
152 D-139 蕹菜・Water spinach. Ipomoea aquatica Forsk.
152 D-140 莧菜・Edible amaranth. Amaranthus mangostanus L.
153 D-141 莧菜・Edible amaranth.
153 D-142 莧菜・Edible amaranth.
154 D-143 芥藍・Kale. Brassica oleracea L. Var. acephala DC.
154 D-144 芥藍・Kale.
154 D-145 芥藍・Kale.
155 D-146 芥藍・Kale.
155 D-147 芥藍・Kale.
155 D-148 黄瓜・Cucumber. Cucumis sativus L.
156 D-149 黄瓜・Cucumber.
156 D-150 黄瓜・Cucumber.
156 D-151 黄瓜・Cucumber.
157 D-152 黄瓜・Cucumber.
157 D-153 黄瓜・Cucumber.
158 D-154 絲瓜・Vegetable sponge. Luffa spp.
158 D-155 絲瓜・Vegetable sponge.
158 D-156 絲瓜・Vegetable sponge.
159 D-157 蕃茄・Tomato. Lycopersicon esculentum Mill.

159 D-158 蕃茄・Tomato.
159 D-159 蕃茄・Tomato.
160 D-160 蕃茄・Tomato.
160 D-161 蕃茄・Tomato.
160 D-162 蕃茄・Tomato.
161 D-163 辣椒・Hot pepper. Capsicum frutescens L.
161 D-164 辣椒・Hot pepper.
162 D-165 菊花・Floret Chrysanthemum. Chrysanthemum morifolium Ramat.
162 D-166 菊花・Floret Chrysanthemum.
162 D-167 菊花・Floret Chrysanthemum.
163 D-168 菊花・Floret Chrysanthemum.
163 D-169 唐菖蒲・Gladioli. Gladiolus hybridus Hort.
164 D-170 玫瑰・Rose. Rosa hybrida Hort.
164 D-171 玫瑰・Rose.
164 D-172 玫瑰・Rose.
165 D-173 玫瑰・Rose.
165 D-174 非洲菊・African daisy. Gerbera jamesonii Bolus.
165 D-175 非洲菊・African daisy.
166 D-176 非洲菊・African daisy.
166 D-177 金露花・Duranta repens.
166 D-178 金露花・Duranta repens.
167 D-179 九重葛・Paper flower. Baugainvillea glabra Choisy.
167 D-180 九重葛・Paper flower.
168 D-181 九重葛・Paper flower.
168 D-182 九重葛・Paper flower.
169 D-183 秋海棠・Wax begonia. Begonia x Semperfiorens-cultorum Hort.
169 D-184 秋海棠・Wax begonia.
169 D-185 秋海棠・Wax begonia.
170 D-186 秋海棠・Wax begonia.
170 D-187 秋海棠・Wax begonia.
170 D-188 秋海棠・Wax begonia.
171 D-189 可可椰子・Coconut. Cocos nucifera L.
171 D-190 可可椰子・Coconut.
171 D-191 可可椰子・Coconut.
172 D-192 袖珍椰子・Parlor palm. chamaedorea elegans Mart.

172 D-193 袖珍椰子・Parlor palm.
172 D-194 袖珍椰子・Parlor palm.
173 D-195 相思樹・Acacia. Acacia sp.
173 D-196 相思樹・Acacia.
173 D-197 相思樹・Acacia.
174 D-198 相思樹・Acacia.
174 D-199 腎蕨・Erect sword fern. Nephrolepis cordifolia K. B. PRESL.

E. 混合氣體污染物・Air Pollutant Combinations.

(二氧化硫＋二氧化氮) (Sulfur Dioxide ＋ Nitrogen Dioxide)

175 E-1 蕃石榴・Guava. Psidium guajava L.
175 E-2 蕃石榴・Guava.
176 E-3 蕃石榴・Guava.
176 E-4 蕃石榴・Guava.
176 E-5 香蕉・Banana. Musa spp.
177 E-6 香蕉・Banana.
177 E-7 香蕉・Banana.
178 E-8 相思樹・Acacia. Acacia sp.

(二氧化硫＋氟化物) (Sulfur Dioxide ＋ Fluorides)

178 E-9 蕃石榴・Guava. Psidium guajava L.
179 E-10 蕃石榴・Guava.
179 E-11 蕃石榴・Guava.
179 E-12 橄欖・Chinese olive. Canarium album Raeusch.
180 E-13 橄欖・Chinese olive.
180 E-14 唐菖蒲・Gladioli. Gladiolus hybridus Hort.
180 E-15 唐菖蒲・Gladioli.
181 E-16 唐菖蒲・Gladioli.
181 E-17 相思樹・Acacia. Acacia sp.
181 E-18 相思樹・Acacia.
182 E-19 相思樹・Acacia.

(臭氧＋氟化物) (Ozone ＋ Fluorides)

182 E-20 木薯・Cassava. Manihot esculenta Crantz.
183 E-21 菸草・Tobacco. Nicotiana tabacum L.
183 E-22 菸草・Tobacco.
183 E-23 菸草・Tobacco.

184	E-24	香蕉・Banana. Musa spp.
184	E-25	香蕉・Banana.
185	E-26	香蕉・Banana.
185	E-27	咖啡・Coffee. Coffea spp.
186	E-28	蓮霧・Wax apple. Syzygium samarangense Merr. et perry.
186	E-29	木瓜・Papaya. Carica papaya Linn.
187	E-30	木瓜・Papaya.
187	E-31	蕹菜・Water spinach. Ipomoea aquatica Forsk.
187	E-32	蕹菜・Water spinach.
188	E-33	茄子・Egg-plant. Solanum melongena L.
188	E-34	茄子・Egg-plant.
188	E-35	茄子・Egg-plant.
189	E-36	黃秋葵・Okra. Hibiscus esculentus L.
189	E-37	黃秋葵・Okra.
189	E-38	黃秋葵・Okra.
190	E-39	毛豆・Vegetable soybean. Glycine Max Merr.
190	E-40	毛豆・Vegetable soybean.
190	E-41	毛豆・Vegetable soybean.
191	E-42	金露花・Duranta repens.
191	E-43	金露花・Duranta repens.
191	E-44	金露花・Duranta repens.

版權所有・不准翻印

All rights reserved. No part of this publication may be reproduced.

空氣污染爲害農作物徵狀診斷圖鑑

Symptom Diagnosis of Air Pollution Injury to Vegetation.

—A Pictorial Atlas—

發行人：杜 金 池

Publisher : Director, C. C. Tu

著 者：林茂盛、張瑞明、陳琳湖

Author : Maw-Sheng Lin, Jui-Ming Chang, Lin-Hu Chen

出版者：臺灣省農業試驗所

臺中縣霧峰鄉中正路189號 電話：(04)3302301

Published by Taiwan Agricultural Research Institute

189 Chung-Cheng Road, Wu-feng 41301, Taichung,

Taiwan, Republic of China. Tel. (04)3302301

承印者：寶成電腦文字排版坊

台中市東區建成路300號 電話：(04)2130862

1994年6月 June, 1994

ISBN 957-00-3844-6

勘 誤 表

Errata List

誤值 (Error)	正確 (Correct)
P 72 B-31 桃. Peach. Amygdalus persica Sieb. et Zucc. 使用品種：新世紀梨 B-32 桃. Peach. B-33 桃. Peach.	P 72 B-31 梨. Pear Pyrus Communis L. 使用品種：新世紀梨 B-32 梨. Pear. B-33 梨. Pear.
P 123 D-58 龍眼. Longan. 使用品種：粉殼 D-59 龍眼. Longan. ~~D-60 龍眼. Longan.~~	P 123 D-58 荔枝. Litchi. Litchi chinensis Sonn. 使用品種：黑葉. D-59 荔枝. Litchi. D-60 荔枝. Litchi.
P 147 D-126 再觀察照片D-126. Again a view of plate D-126.	P147 D-126 再觀察照片D-125. Again a view of plate D-125.
P 197 72 B-31 桃 • Peach. Amygdalus persica Sieb. et Zucc. 72 B-32 桃. Peach. 72 B-32 桃. Peach.	P 197 72 B-31梨. Pear Pyrus Communis L. 72 B-32 梨. Pear. 72 B-33 梨. Pear.
P 201 123 D-58 龍眼. Longan. 123 D-59 龍眼. Longan. 123 D-60 龍眼. Longan.	P 201 123 D-58 荔枝. Litchi. Litchi chinensis Sonn. 123 D-59 荔枝. Litchi. 123 D-60 荔枝. Litchi.

統 一 編 號
0308598300020
ISBN 957-00-3844-6